

5 ALFALFA HYBRIDS HAVING AT LEAST 75% HYBRIDITY

CROSS-REFERENCE TO RELATED APPLICATIONS

Not applicable.

BACKGROUND OF THE INVENTION

10

Alfalfa (*Medicago sativa*) has often been referred to as the "Queen of Forages" because it is an excellent source of protein and digestible fiber, and because its wide adaptation. Alfalfa is highly effective in nitrogen fixation, and is frequently planted in crop rotation to replenish nutrients depleted from the soil by other crops
15 such as corn.

15

Efforts to develop alfalfa varieties having improved traits and increased production have focused on breeding for disease, insect, or nematode resistance, persistence, adaptation to specific environments, increased yield, and improved quality. Breeders have had less success in breeding for yield and quality per se,
20 although methods directed toward increasing herbage quality and forage yield have been developed. For example, see U.S. Pat. No. 4,045,912, which is incorporated herein by reference.

20

Breeding programs typically emphasize maximizing heterogeneity of a given alfalfa variety to improve yield and stability. However, this generally results in wide
25 variations in characteristics such as flowering dates, flowering frequency, development rate, growth rate, fall dormancy and winter hardiness. Prior art breeding methods do not emphasize improving the uniformity of these characteristics.

25

An important economic consideration in the development of hybrid alfalfa strains is the ability of such hybrids to produce acceptable yields of alfalfa seed. U.S.
30 Patent No. 4,045,912, incorporated by reference herein, discloses a process for producing alfalfa seed by random pollination of male sterile plants (produced by a cross of a cytoplasmic male sterile line and maintainer alfalfa line) and a male fertile line, with the ratio of male sterile plants to male fertile plants in the range of from 1:1 to 3:1.

30

There is a need in the art for producing alfalfa hybrids having agronomically desirable traits and breeding methods that result in a high degree of hybridity, uniformity of selected traits, and acceptable seed yields.

5

BRIEF SUMMARY OF THE INVENTION

6401 - In one aspect, the present invention provides a *Medicago sativa* hybrid seed or cultivated alfalfa seed designated DS9705Hyb ((A833 x B209) x (Thor, DS9761 and C580)), and deposited under the terms of the Budapest Treaty on December 4, 2000 with the America Type Culture Collection (ATCC), Manassas, VA, under Accession
10 Number XXXXX.

The present invention includes a *Medicago sativa* hybrid plant or cultivated alfalfa plant derived from the seed deposited under Accession Number XXXXX. The plant may be grown directly from the seed deposited under Accession Number XXXXX, or may be obtained indirectly from a plant grown directly from the seed by
15 any suitable means. For example, the plant may be generated from seed produced by a plant grown directly from the seed, from a cutting taken from a plant grown directly from the seed, or from tissue culture or callous derived from cells from a plant grown directly from the seed. The invention includes succeeding generations of plants derived from plants grown from the seed of Accession Number XXXXX.

20 In other aspects, the present invention includes the pollen and ovule of a plant derived from the seed deposited under Accession Number XXXXX.

In another aspect, the present invention provides a method of producing alfalfa seeds having at least 75% hybridity comprising the steps of: crossing cytoplasmic male sterile alfalfa plants with maintainer alfalfa plants to produce cytoplasmic male
25 sterile hybrid seed; selectively harvesting seed from the cytoplasmic male sterile plants; crossing plants grown from the seed of male sterile hybrid plants by male fertile alfalfa plants by pollination by growing the seed from male sterile hybrid plants with seed from at least one male fertile alfalfa plant, the male sterile and male fertile seed planted at a ratio of about four male sterile seeds to every one male fertile seed;
30 and non-selectively recovering seeds from the pollinated alfalfa plants.

Another aspect of the invention provides seed comprising at least 75% hybrid seed, the seed produced by crossing cytoplasmic male sterile alfalfa plants with maintainer alfalfa plants to produce cytoplasmic male sterile hybrid seed; selectively harvesting seed from the cytoplasmic male sterile plants; crossing plants grown from

the seed of male sterile hybrid plants by male fertile alfalfa plants by pollination by growing the seed from male sterile hybrid plants with seed from at least one male fertile alfalfa plant, the male sterile and male fertile seed planted at a ratio of about four male sterile seeds to every one male fertile seed; and non-selectively recovering
5 seeds from the pollinated alfalfa plants.

Optionally, the percentage hybridity can be verified using either genetic or morphological markers.

It is an advantage of the present invention that the method of producing hybrid alfalfa seed results in seed with at least 75% hybridity in commercially acceptable
10 yields.

It is an advantage that any cytoplasmic male sterile alfalfa plants and maintainer alfalfa plants may be selected for use in the practice of the present invention to obtain a cytoplasmic male sterile hybrid plant.

It is a further advantage of the present invention that the forage yields of fields
15 planted with 75% hybrid alfalfa are comparable to forage yields of fields planted with 100% hybrid seed, a feature that allows for high forage yields and high seed production.

Other features and advantages of the invention will be apparent upon review of the specification.
20

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

Not applicable.

DETAILED DESCRIPTION OF THE INVENTION

25 In one aspect, the present invention includes the hybrid alfalfa seed deposited under the Budapest Treaty on December 4, 2000 with the American Type Culture Collection in Manassas, VA as Accession Number XXXXX, and plants or plant parts derived from the seed deposited as Accession Number XXXXX.

By "a plant derived from the seed deposited as Accession Number XXXXX",
30 it is meant a plant that is grown directly from the seed deposited as Accession Number XXXXX, or a plant that is obtained indirectly from a plant grown from the seed deposited as Accession Number XXXXX. Plants obtained indirectly from a plant grown from the seed deposited as Accession Number XXXXX may be grown from a vegetative cutting, seed, callus or tissue culture obtained from a plant or plant

part grown from the seed deposited as Accession Number XXXXX or a clonal plant thereof.

5 The present invention also provides a method of obtaining hybrid alfalfa lines using cytoplasmic male sterile alfalfa lines (A lines), maintainer alfalfa lines (B lines), and male fertile pollenizer lines (C lines) as described in detail in the examples.

Male sterile A lines may be identified by evaluating pollen production using the Pollen Production Index (P.P.I.), which recognizes four distinct classes:

1. Male Sterile Plants (MS) PPI=0

10 No visible pollen can be observed with the naked eye when flower is tripped with a black knife blade.

2. Partial Male Sterile Plant (PMS) PPI=0.1

A trace of pollen is found with the naked eye when flower is tripped with a black knife blade.

3. Partial Fertile Plant (PF) PPI=0.6

15 Less than a normal amount of pollen can be observed with the naked eye when flower is tripped with a black knife blade.

4. Fertile Plant (F) PPI=1.0

Normal amounts of pollen can be observed when flower is tripped with a black a knife blade.

20 The cells of the cytoplasmic male sterile (A line) alfalfa plants contain sterile cytoplasm and the non-restorer gene. The maintainer line (B line) is a male and female fertile plant, and when crossed with an A line plant, maintains the male sterility of the cytoplasmic male sterile plant in the progeny. The cells of a maintainer line plant contain normal cytoplasm and the non-restorer gene. Methods for
25 identifying cytoplasmic male sterile and maintainer lines of alfalfa are well known to those versed in the art of alfalfa plant breeding (e.g., see U.S. Patent No. 3,570,181, which is incorporated by reference herein). A pollenizer line (C line) is a fertile plant containing both male and female parts.

Briefly, the method of the invention is performed as follows:

30 1. Alfalfa plants with desirable agronomic traits are selected. Male sterile A line plants are selected from male sterile ("female") populations, maintainer B line plants are selected from maintainer populations, and pollenizer C line plants are selected from restorer populations, or from clonal or synthetic populations.

2. The selected A and B lines are grown from cuttings or seed and cross pollinated using bees to produce hybrid male sterile breeder and foundation seeds. Seeds are harvested from cytoplasmic male sterile plants only.

3. Selected pollenizer plants are selfed or interpollinated by bees to produce
5 breeder and foundation pollenizer seeds and the seed is harvested in bulk.

4. For large scale commercial production of hybrids, male sterile seeds and pollenizer seeds are planted at a ratio of male sterile seeds and male fertile (pollenizer) seeds of about 4:1, and the plants grown therefrom are pollinated.

5. Seeds are harvested in bulk from the plants grown from the seed of step 4,
10 above.

6. Optionally, the percentage hybridity can be determined using either genetic or morphological markers.

Cytoplasmic male sterile lines may be maintained by vegetative cuttings. Maintainer lines can be maintained by cuttings or self-pollination. Male sterile
15 hybrids can be obtained by cross-pollinating cytoplasmic male sterile plants with maintainer plants. Pollenizer lines can be maintained by selfing or, if more than two clones are used, by cross-pollination.

Preferably, at least one of the alfalfa plant lines used in developing alfalfa hybrids according to the method of the present invention has at least one desirable
20 agronomic trait, which may include, for example, resistance to disease or insects, cold tolerance, increased persistence, greater forage yield or seed yield, improved forage quality, uniformity of growth rate, and uniformity of time of maturity.

In the controlled pollination step, the cytoplasmic male sterile plants are typically grown in separate rows from the maintainer plants. The plants are pollinated
25 by pollen-carrying insects, such as bees. Segregating the male sterile and maintainer plants facilitates selective harvest of hybrid seed from the cytoplasmic male sterile plants.

The male sterile seed and male fertile seed is preferably provided as a random mixture of the seed in a ratio of about 4:1, which would provide for random
30 distribution of the male sterile and male fertile plants grown therefrom and random pollination of the alfalfa plants. As one of skill in the art will appreciate, one could also practice the method of the invention using designed distribution of male sterile hybrid and male fertile lines within a field and subsequent pollination by pollen-carrying insects.

In the examples below, male sterile line A833, maintainer line B209, and pollenizer lines Thor, DS9671, and C580 were used. One of ordinary skill in the art will appreciate that any suitable male sterile line, maintainer line, and pollenizer line could be successfully employed in the practice of the method of the invention.

5 The term "hybrid" as used herein is intended to refer to a cross between a male sterile line and a pollenizer line in which at least about 75% of the seeds differ genetically from the parent lines. The term S1 refers to the first generation of selfing.

 Preferably, the method of the invention produces seed having at least about 75% hybridity (i.e., at least 75% of the seeds differ genetically from the male and
10 female parent lines). Estimates of hybridity showed that the method of the invention yielded seeds comprising about 90% hybrids. However, forage yields obtained from fields seeded with 75 or 80% hybrid seed were comparable to forage yields obtained from fields planted with 100% hybrid seeds. Therefore, it is expected that hybrid
15 production seed having 75% hybrid seeds would give forage yields comparable to those obtained from fields planted with 100% hybrid seed.

 Preferably, the seed produced by the method of the invention is produced in commercially acceptable yields. Of course, what constitutes a "commercially acceptable seed yield" may vary from line to line, depending on the totality of hybrid's characteristics. Preferably, seed production of plants grown from male
20 sterile hybrid seed crossed with plants grown from male fertile seed planted in a ratio of 4:1 is at least 80% of the average seed production obtained from selfing the male fertile parent. More preferably still, seed production is at least 85% as great as, or even as much as 95% or 100% as great as seed production by the male fertile parent. What constitutes a commercially acceptable seed yield may be made with reference to
25 typical seed production of synthetic alfalfa varieties. In the examples below, hybrid alfalfa seed production yields were comparable to those obtained from synthetic alfalfa varieties such as Magnum V or Magnum IV.

 The following non-limiting examples are intended to be purely illustrative.

EXAMPLES

Scoring of plant size

Two-month-old seedlings started in the greenhouse were transplanted to the field in April of 1999 with a spacing of 40 inches between plants. Plants were trimmed off in August of 1999 and herbage regrowth visually scored in September of the same year. Plants were scored from 1 to 9, with "1" being the least herbage and "9" greatest herbage. The mean, variance and range of visual scores of plant size are provided in Table 1.

Upon visual inspection, hybrids A833 x B209 and A833 x B209 (S1) appeared more vigorous than the synthetic varieties Vernal and Saranac and more vigorous than the (S1) progeny of selected clones (Thor, DS9671, C580 and B209). The hybrids also show greater uniformity in size than synthetic varieties, as is evidence by the lower variance.

Survival of over wintered plants

The percentage of plants that survived over the winter of 1999 was determined (Table 2). Hybrids A833 x 209 and A833 x 209 (S1) had the highest survival rates at 95.8% and 95.0%, respectively. The synthetic varieties Vernal and Saranac, which had survival rates of 82.2% and 87.9%, respectively.

The survival rate of the progeny of different individual clones varied from 80.4 to 92.7%. Hybrid A833 x B209 showed remarkable improvement in winter survival over (S1) progeny of the male parent B209. The formation of hybrids can enhance the ability to survive winter stress.

Assessment of spring vigor

Spring vigor of alfalfa plants was assessed and scored from 1-9, with "1" being less vigorous and "9" being most vigorous. The results are summarized in Table 3.

Hybrids, A833 x B209 and A833 x B209 (S1) showed greater spring vigor and uniformity relative to the synthetic varieties Vernal and Saranac, as well as greater vigor and uniformity than the (S1) progeny of selected clones (Thor, DS9671, C580 and B209).

Flowering dates

The first and second Julian flowering dates were recorded in June and July-August, respectively, for synthetic varieties Vernal and Saranac, (S1) progeny of

selected clones (Thor, DS9761, C580 and maintainer line B209), and hybrids A833 x B209 and A833 x B209 (S1). The Julian flowering date is measured in days from January 1st and is expressed in days. The results are presented in Table 4 and 5.

The hybrids A833 x B209 and A833 x B209 (S1) had more uniform flowering dates than progeny of the pollinators, maintainer line, or synthetics, as is evidenced by a relatively low variance.

Flower Color

Flower colors were recorded for each alfalfa plant (Table 6).

Pollen Production Index

Four male fertility classifications, male sterile (MS), partial male sterile (PMS), partial fertile (PF), and fertile (F) were used in this research project. The distribution for male sterility of hybrids, A833 x B209 and A833 x B209 (S1) were recorded and the results are presented in Table 7.

P.P.I. for A833 x B209 and A833 x B209 (S1) are .077 and .086 respectively. Slightly increased P.P.I. values were obtained when S1 progeny were used as maintainer line.

Disease resistance of hybrid and hybrid components

The response of alfalfa plant hybrid DS9705Hyb ((A833 x B209) x (Thor, DS9761 and C580)) and hybrid components (B209, A833 x B209, Thor, DS9761 and C580) to various diseases was evaluated according to the "Standard Tests to Characterize Alfalfa Cultivars", approved by the North American Alfalfa Improvement Conference. The resistance or susceptibility of the hybrid or hybrid components to anthracnose, bacterial wilt, Fusarium wilt, Verticillium wilt, Phytophthora root rot, Aphanomyces root rot (Race 1), stem nematode and root knot nematode was assessed. For each disease tested, appropriate check cultivars, including resistant and susceptible cultivars, were employed as controls. The results are presented in Tables 8 through 23.

For each type of disease tested, each line of plants was assigned to one of five classes of resistance according to the percentage of resistant plants as follows:

	<u>Class</u>	<u>% Resistant plants</u>
30	Susceptible	<6
	Low resistant	6-14
	Moderately resistant	15-30
	Resistant	31-50
35	Highly resistant	>50

DS9705Hyb was found to be highly resistant to bacterial wilt, Fusarium wilt, Phytophthora root rot, root knot nematode; resistant to anthracnose (race 1), Verticillium wilt, stem nematode and moderately resistant to Aphanomyces root rot (Race 1).

5 Forage yields

Forage yields of hybrid DS9705Hyb and the hybrid components B209, A833 x B209, Thor, DS9761 and C580 were measured and are presented on Tables 24 and 25. The hybrid DS9705Hyb showed average yields that were 12% higher than those obtained with Vernal over eighty five harvests and twelve test locations.

10 Persistence Advantage of the Hybrids

Persistence of hybrid DS9705Hyb showed a 15% advantage over check variety Vernal in five test environments. (Table 26)

Seed yield of hybrid and hybrid components

15 Alfalfa seed yield is usually greatly affected by cultural management, field conditions and year. Three fields of the alfalfa hybrid DS9705Hyb were planted in the spring of 2000. Hybrid alfalfa seed was obtained at a yield of about 400, 500, and 600 pounds per acre, a yield comparable to that of synthetic varieties. The synthetic variety Magnum V produced seed at a rate of about 400, 500, and 600 pounds per acre; Magnum IV yielded about 375, 450, and 550 pounds seed/per acre in the same
20 year. Seed yield of male sterile foundation seeds A833 x B209 was significantly lower than maintainer line B209.

Percentage of hybrid required for maximum forage yield

25 The hybrid A19-616 x Thor, showed a forage yield 16% higher than that of the inbred AP-51. To determine the hybrid percentage required for maximum forage yield, fields were seeded with mixtures of hybrid and inbred seeds in different ratios separated by 10% intervals, ranging from 100% hybrid and 0% inbred to 0% hybrid and 100% inbred. The results of the forage yield experiments, which were conducted in Clinton, WI, Arlington, WI, and Rosemount, MN in 1997 and 1998, are presented in Table 27.

30 Seed comprising 100% hybrid A19-616 x Thor yielded 6.0 tons/per acre, whereas inbred AP-51 yielded 5.19 tons/per acre, with a L.S.D.(.05) of 0.2. The results show no difference in forage yields of fields planted with from 80-100% hybrid seed, and a slight decrease in forage yields of fields planted with from 50-70% hybrids.

Percent hybridity of the hybrid DS 9705Hyb

Seed obtained from non-selective harvest of male sterile hybrids crossed by male fertile plants planted at a ratio of 4:1 includes both hybrid and non-hybrid seed. Molecular markers were identified to allow differentiation of hybrids from parental lines as follows.

Parental plants B209, DS9671, C580 and Thor were selfed and pure S1 seed was generated. Amplified fragment length polymorphism (AFLP) analysis of DNA isolated from plants derived from the parental seed of B209, DS9671, C580, Thor; the hybrid seed of A833xB209; and hybrid seed of DS9705Hyb was conducted by KeyGene Molecular Marker Services (P.O. Box 216, 6700 AE Wageningen, The Netherlands). The seeds were germinated and grown into plants. DNA was isolated from a pool of 20 individuals from each pedigree. The parental samples were subjected to AFLP using 15 primer combinations, which identified polymorphic markers between the B209, Thor, C580, DS9671 and A833xB209 parents. The two primer pairs, designated E32/M49 and E36/M50, were found to identify the greatest numbers of polymorphisms and were used to amplify DNA in production lots of DS9705Hyb. Primers E32, M49, E36, and M50 have the sequences shown in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, and SEQ ID NO:4, respectively. Ninety plants were grown from DS 9705Hyb production lot seed and DNA was isolated from each plant individually. The DNA was subjected to AFLP using the two primer pairs. Of the 90 plants tested, 90% were hybrid plants and 10% were male selfs.

Table 1 Herbage regrowth scores of alfalfa plants.

Herbage regrowth					
		Number	Mean	Variance	Range
5					
	Vernal	136	4.35	1.07	1-7
	Saranac	155	4.38	0.95	2-7
	Thor	129	4.37	0.69	1-7
10	DS9761	115	4.43	0.51	3-6
	C580	175	4.26	0.61	1-6
	B209	200	4.34	0.57	2-7
	A833x209	116	4.71	0.57	2-7
	A833x209(S1)	114	4.61	0.72	1-6
15	Planted: 4/23/99 at the Dairyland Research Station near Clinton, WI				
	Scored: 9/99				
	Rating scale: 1 is least amount of herbage				
	9 is greatest amount of herbage				

20

Table 2 Survival of over wintered alfalfa plants

		<u>Total plants</u>	<u>Live</u>	<u>Dead</u>	<u>Survival (%)</u>
25	Vernal	163	134	29	82.2
	Saranac	174	153	21	87.9
	Thor	150	131	19	87.3
	DS9671	124	115	9	92.7
30	C580	202	177	25	87.6
	B209	250	201	49	80.4
	A833x209	120	115	5	95.8
	A833x209 (S1)	120	114	6	95.0

35 Planted: 4/23/99 at the Dairyland Research Station near Clinton, WI
 Scored: 5/00
 Rating scale: By "live" is meant an actively growing plant
 By "dead" is meant a non-growing plant

40

Table 3 Spring vigor of alfalfa plants

		<u>Number</u>	<u>Mean</u>	<u>Variance</u>	<u>Range</u>
5	Vernal	164	5.17	6.82	1-8
	Saranac	174	5.81	5.57	1-9
	Thor	150	5.73	5.51	1-8
	DS9671	124	5.89	3.45	1-9
10	C580	202	5.59	5.98	1-9
	B209	250	5.28	5.41	1-9
	A833x209	120	6.88	2.59	1-9
	A833x209 (S1)	120	6.40	2.93	1-8

15 Planted: 4/23/99 at the Dairyland Research Station near Clinton, WI
 Scored: 5/00
 Rating Scale: 1 is least vigorous; 9 is most vigorous

20 **Table 4** First Julian flowering dates (expressed in days from January 1 to first flowering)

		<u>Number</u>	<u>Mean</u>	<u>Variance</u>	<u>Range</u>
	Vernal	133	148.6	31.1	135-159
	Saranac	154	148.4	28.31	134-159
25	Thor	129	148.5	30.57	115-159
	DS9671	115	148.4	20.38	142-159
	C580	176	147.7	19.50	136-159
	B209	201	147.4	18.01	135-159
	A833x209	115	145.0	11.74	137-157
30	A833x209 (S1)	114	147.2	23.70	135-159

Planted: 4/23/99 at the Dairyland Research Station near Clinton, WI
 Data collected in 2000 growing season

35 **Table 5** Second Julian flowering dates (expressed in days from January 1 to second flowering)

		<u>Number</u>	<u>Mean</u>	<u>Variance</u>	<u>Range</u>
40	Vernal	133	186.5	12.87	177-195
	Saranac	152	185.8	11.67	175-195
	Thor	132	186.2	11.41	177-195
	DS9761	115	187.1	8.19	180-195
	C580	177	185.6	9.41	180-195
45	B209	201	184.5	13.21	176-195
	A833x209	115	185.5	4.69	177-195
	A833x209 (S1)	114	185.9	7.58	177-195

50 Planted: 4 23 99 at the Dairyland Research Station near Clinton, WI
 Data collected in 2000 growing season

Table 6 Flower color classification

	<u>N</u>	<u>P</u>	<u>BP</u>	<u>B</u>	<u>G</u>	<u>V</u>	<u>W</u>
5							
	Vernal	126	119	0	0	0	3
	Saranac	149	149				4
	Thor	129	128				1
10	DS9761	113	112		1		
	C580	174	172	1	1		
	B209	194	194				
	A833x209	114	114				
15	A833x209 (S1)	113	113				

P=purple; BP=blue purple; B=blue, G=green; V=variegated; and W=white.

20

Table 7 Pollen production of readings of A833xB209 and A833 x B209 (S1) in June 2000 at Clinton, Wisconsin

	<u>N</u>	<u>Fertility classifications</u>				
25		<u>MS</u>	<u>PMS</u>	<u>PF</u>	<u>F</u>	<u>P.P.I</u>
	A833xB209 114	91	14	4	5	.077
	A833xB209 (S1) 106	89	7	4	6	.086
30		<u>Rating</u>	<u>PPI Value</u>			
	Fertility classifications:	MS	0.0			
		PMS	0.1			
		PF	0.6			
35		F	1.0			

Table 8 Resistance to anthracnose (Race 1) (*Colletotrichum trifolii*)

Test conducted by Dairyland Research at Clinton, WI

5	Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
10	B209	R	1996	1	29.3	30.5
	A833xB209	R		1	33.3	34.7
	C580	R		2	33.4	34.8
	DS9671	HR		2	54.3	56.6
15	Thor	MR		2	28.0	29.2
	Check Varieties					
	1. Saranac AR	R			43.2	45.0
	2. Saranac	S			5.0	5.2
20	Test Mean:				43.8	45.6
	L.S.D. (.05%)				5.2	
	C.V.(%)				8.6	
25	Test Conducted in Lab					

Note: Unadjusted % R is the actual raw data summary.

Adjusted % R is transformed to the standards of the resistant check.

Table 9 Resistance to anthracnose (Race 1) (*Colletotrichum trifolii*)

Test conducted by Dairyland Research at Clinton, Wisconsin

5	Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
10	DS9705Hyb	R	1999	1	39.2	43.9
	Check Varieties					
	1. Saranac	ARR			40.2	45.0
15	2. Saranac	S			3.2	3.6
	Test Mean:				32.1	42.1
	L.S.D. (.05%)				6.8	
20	C.V. (%)				11.6	
	Test conducted in Lab					

Table 10 Resistance to bacterial wilt (*Clavibacter michiganense*)

Test conducted by Dairyland Research at Clinton, WI.

25	Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R	Score A.S.I.
30	B209	HR	1998	1	63.0	73.5	1.40
	A833xB209	HR		1	79.4	86.3	1.37
	C580	HR		2	56.2	65.6	1.96
	DS9671	HR		2	48.0	56.0	2.25
	Thor	HR		2	54.0	63.0	2.28
35	Check Varieties:						
	1. Vernal	R			36.0	42.0	1.80
	2. Narragansett	S			3.0	3.5	3.8
40	Test Mean:				61.0	71.1	2.4
	L.S.D. (.05%)				13.0		.21
	C.V.(%)				30.3		13.8
45	Test conducted in field.						

Note: Score A.S.I. is the average severity index. Ratings range from 0-5 with a score of "0" indicating no discernable disease symptoms and "5" indicating plant death.

Table 11 Resistance to bacterial wilt (*Clavibacter michiganense*)

Test conducted by Dairyland Research at Clinton, Wisconsin

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R	Score A.S.I.
DS9705Hyb	HR	1999	1	63.5	74.1	1.6
Check Varieties						
1. Vernal	R			46.0	53.6	1.8
2. Narragansett	S			3.0	3.5	3.8
Test Mean:	51.0			59.5	2.20	
L.S.D. (.05%)				8.2		.35
C.V. (%)				14.2		9.60
Test conducted in field						

Table 12 Resistance to Fusarium wilt (*Fusarium oxysporum*)

Test conducted by Dairyland Research at Clinton, WI

[illegible]

Table 13 Resistance to Fusarium wilt (*Fusarium oxysporum*)

Test conducted by Dairyland Research at Clinton, WI

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R	Score or A.S.I.
DS9705Hyb	HR	1999	1	72.7	78.5	1.42
Check Varieties						
1. Agate	R			50.0	54.0	1.56
2. MNGN-1	S			0.0	0.0	3.20
Test Mean:				62.9	67.9	1.47
L.S.D. (.05%)				5.3		.26
C.V.(%)				9.85		6.25
Test conducted in field.						

Table 14 Resistance to Verticillium wilt (*Verticillium albo-atrum*)

Test conducted by Dairyland Research at Clinton, Wisconsin

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
B209	R	1998	1	37.0	32.9
A833xB209	R		1	43.0	38.2
C580	R		2	38.5	34.2
DS9671	R		2	35.0	31.6
Thor	R		2	41.0	36.4
Check Varieties					
1. Vertus	R			45.0	40.0
2. Saranac	S			4.8	4.3
Test Mean:				38.0	33.8
L.S.D.(05%)				7.8	
C.V.(%)				13.1	
Test conducted in Lab					

Table 15 Resistance to Verticillium wilt (*Verticillium albo-atrum*)

Test conducted by Dairyland Research at Clinton, WI

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
DS9705Hyb	R	1999	1	36.8	33.2
Check Varieties					
1. Vertus	R			44.3	40.0
2. Saranac	S			5.0	4.5
Test Mean:				22.8	20.6
L.S.D. (.05%)				9.9	
C.V. (%)				16.2	
Test conducted in Lab					

Table 16 Resistance to Phytophthora root rot (*Phytophthora medicaginis*)

Test conducted by Dairyland Research at Clinton, Wisconsin.

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
B209	MR	1996	1	21.8	22.8
A833xB209	R		1	38.5	40.3
C580	R		2	42.1	44.1
DS9671	HR		2	56.5	59.2
Thor	HR		2	66.2	69.4
Check Varieties					
1. Agate	R			31.5	33.0
2. Saranac	S			0.0	0.0
Test Mean:				12.5	13.1
L.S.D. (.05%)				8.6	
C.V. (%)				14.8	
Test conducted in the Lab.					

Table 17 Resistance to Phytophthora root rot (Phytophthora medicaginis)

Test conducted by Dairyland Research at Clinton, Wisconsin.

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
DS9705Hyb	HR	1999	1	59.2	60.0
Check Varieties					
1. Agate	R			32.6	33.0
2. Saranac	S			0.0	0.0
Test Mean:				48.6	49.2
L.S.D. (05%)				10.6	
C.V. (%)				16.8	
Test conducted in the Lab.					

Table 18 Resistance to stem nematode (*Ditylenchus dipsaci*)

Test conducted by Dairyland Research at Clinton, Wisconsin.

25

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R	Score A.S.I.
30	B209	R	1996	1	39.0	42.2
	A833xB209	R		1	35.0	37.8
	C580	R		2	33.2	35.9
	DS9671	R		2	34.0	36.8
	Thor	R		2	29.0	31.4
35	Check Varieties					
	1. Lahontan	R			37.0	40.0
	2. Saranac	S			1.0	1.1
40	Test Mean:				29.0	31.4
	L.S.D. (05%)				7.6	
	C.V. (%)				11.6	
	Test conducted in the Lab.					

45

Table 19 Resistance to stem nematode (*Ditylenchus dipsaci*)

Test conducted by Dairyland Research at Clinton, Wisconsin.

5	Variety	Resistance Class	Year	Syn Tested	Unadjusted Gen % R	Adjusted % R	Score
	A.S.I.						
10	DS9705Hyb	R	1999	1	35.0	37.8	3.13
	Check Varieties						
	1. Lahontan	R			37.0	40.0	3.22
	2. Saranac	S			1.0	1.1	4.16
15	Test Mean:				29.0	31.4	3.6
	L.S.D.				7.6		0.36
	C.V. (%)				11.6		7.6
	Test conducted in the Lab.						
20							

Table 20 Resistance to Aphanomyces root rot (Race 1) (*Aphanomyces euteiches*)

Test conducted by Dairyland Research at Clinton, Wisconsin.

25	Variety	Resistance Class	Year	Syn Tested	Unadjusted % R	Adjusted % R
	B209	LR	1996	1	11.2	9.5
30	A833xB209	MR		1	20.0	17.0
	C580	MR		2	26.0	22.1
	DS9671	MR		2	19.2	16.3
	Thor	MR		2	23.8	20.2
35	Check Varieties					
	1. WAPH-1	R			58.9	50.0
	2. Saranac	S			0.0	0.0
40	Test Mean:				12.3	10.4
	L.S.D. (.05%)				6.8	
	C.V. (%)				9.9	
	Test conducted in the Lab					

Table 21 Resistance to Aphanomyces root rot (Race 1) (Aphanomyces euteiches)

Test conducted by Dairyland Research at Clinton, Wisconsin.

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
DS9705Hyb	MR	1999	1	25.5	26.2
Check Varieties					
1. WAPH-1	R			48.7	50.0
2. Saranac	S			0.0	0.0
Test Mean:				15.6	16.0
L.S.D. (.05%)				9.8	
C.V. (%)				12.6	
Test conducted in the Lab.					

Table 22 Resistance to Root-knot nematode (Meloidogyne halpa)

Test conducted by Dairyland Research at Clinton, Wisconsin.

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
B209	R	1996	1	49.0	49.6
A833xB209	HR		1	57.0	57.6
C580	R		2	42.2	42.5
DS9671	HR		2	59.0	59.6
Thor	HR		2	56.0	56.6
Check Varieties					
1. Nev. SynXXHR				89.0	90.0
2. Lahontan	S			7.0	7.1
Test Mean:				61.0	61.7
L.S.D.(.05%)				8.6	
C.V. (%)				13.6	
Test conducted in the Lab.					

Table 23 Resistance to Root-knot nematode (*Meloidogyne halpa*)

Test conducted by Dairyland Research at Clinton, Wisconsin.

5

Variety	Resistance Class	Year Tested	Syn Gen	Unadjusted % R	Adjusted % R
---------	------------------	-------------	---------	----------------	--------------

10

DS9705Hyb HR 1999 1 58.2 58.9

Check Varieties

1. Nev. Syn XXHR 89.0 90.0

2. Lahontan S 7.0 7.1

15

Test Mean: 58.0 58.6

L.S.D. (.05%) 9.2

C.V. (%) 16.5

Test conducted in the Lab.

20

Table 24 Forage Yield of Hybrid DS9705Hyb and Vernal Alfalfa

	Variety		Harvests
	DS9705Hyb Yield (tons/acre)	Vernal Yield (tons/acre)	
FIELD SITE/YEAR			
Lansing, MI 1997-99	13.63	10.88	8
Rosemount, MN 1997-99	12.49	11.48	6
Ames, IA 1997-99	15.54	12.98	8
Arlington, WI 1997-99	11.08	9.48	8
Marshfield, WI 1997-99	11.33	10.01	8
Lancaster, WI 1997-99	12.09	10.53	8
Clinton, WI 1997-99 Early	13.06	12.37	8
Clinton, WI 1997-99 Medium	12.64	11.72	8
Clinton, WI 1997-99 Late	13.69	12.61	6
Clinton, WI 1998-99 Early	9.8	8.8	6
Clinton, WI 1998-99 Medium	9.2	8.5	6
Clinton, WI 1998-99 Late	10.6	9.7	5
Total	145.15	130.06	85
Percent advantage	12%	-	

25

Note: Clinton, WI locations are group by maturity at harvest. Early harvests were conducted about 7 days earlier than medium harvests, and late harvests were conducted about 5 days later than medium harvests.

Table 25 Forage Yield Data

Data collected across at various Midwest locations

Planting years: 94-98

5 Harvest years: 94-99

	<u>Component</u>	<u>Percent Vernal</u>	<u>Total Tons</u>	<u>Number of Harvests</u>
10	A833xB209	112.5	235.5	186
	B209	101.8	126.2	86
	DS9671	110.8	141.86	124
15	Thor	112.9	60.62	32
	C580	114.9	13.62	8
20	<u>Hybrid</u>			
	DS9705Hyb	112	145.15	85

Table 26 Persistence (% stand) of alfalfa DS9705Hyb and synthetic alfalfa varieties.

25

	Early (% stand)	Medium (% stand)	Late (% stand)	Appleton, WI (% stand)	Manawa, WI (% stand)	Mean	Advantage 1
DS9705Hyb	83	82	83	88	87	84.6	115
Magnum IV	78	79	81	83	82	80.6	
Magnum III- Wet	72	77	77	83	82	80.6	
DK127	73	78	77	77	83	77.6	
Legendairy 2.0	72	75	77	82	85	78.2	
P5312	77	80	82	82	82	80.6	
P5454	77	82	80	83	85	81.4	
Vernal	73	77	77	68	73	73.6	100

1. Advantage= (mean % stand for DS9705 / mean % stand for Vernal) x 100

Early, medium and late locations are at the Dairyland Research Station at Clinton, WI, planted 5/97 and scored 5/2000.

30 Appleton, WI, planted 5/97 and scored:10/2000.

Manawa, WI, planted 5/97 and scored 5/99.

Table 27 Forage yield as a function of percent hybrids planted in May, 1996 in Clinton, WI., Arlington, WI., and Rosemount, MN. and harvested in 1997 and 1998.

5	<u>Pedigree</u>	
	Hybrid:	A19-616xThor
	Male:	AP-51
10	<i>Ratio Of:</i>	
	<u>HYBRID</u>	<u>INBRED</u>
		<u>YIELD (TONS/ACRE)</u>
	100	0
15	90	10
	80	20
	70	30
	60	40
	50	50
20	40	60
	30	70
	20	80
	10	90
	0	100
25	LSD(.05)	0.2
	CV(%)	6.11
30		
	Number of harvests	<u>1997</u>
		<u>1998</u>
35	Location:	
	Arlington, WI	4
	Rosemount, MN	3
	Clinton, WI	4
40		

The present invention is not limited to the exemplified embodiments, but is intended to encompass all such modifications and variations as come within the scope of the following claims.